



Review Article

# Phytochemical Diversity and Multispectral Pharmacological Activities of *Azadirachta indica* A.Juss. (Meliaceae): An Evidence-Based Review of Bioactive Constituents, Therapeutic Mechanisms, and Emerging Applications

Archit Kapil<sup>1</sup>, Anshul Anshul<sup>1</sup>, Vansh Gupta<sup>1</sup>, Sandeep Kumar Barwal<sup>1,\*</sup> and Harsh Singh<sup>1</sup>

<sup>1</sup> School of Botany, Maa Shakumbhari University, Punwarka, Saharanpur 247120, Uttar Pradesh, India

\* Corresponding author: [barwal.sk18@gmail.com](mailto:barwal.sk18@gmail.com)

## Abstract

*Azadirachta indica* A.Juss. (Meliaceae), commonly designated as neem, is one of the most extensively investigated medicinal trees in tropical and subtropical phytomedicine. Its remarkable pharmacological versatility stems from an exceptionally diverse phytochemical arsenal distributed across all anatomical parts, including leaves, bark, seeds, flowers, and roots. The principal bioactive constituents belong to two broad chemical categories: isoprenoid compounds most notably the tetra-nortri-terpenoid azadirachtin, nimbin, nimbidin, and salannin and non-isoprenoid compounds encompassing flavonoids, tannins, alkaloids, saponins, coumarins, and phenolic glycosides. Collectively, these constituents confer upon the plant a broad spectrum of pharmacological properties that have been substantiated by contemporary experimental investigations, including antimicrobial, antioxidant, immunomodulatory, neuroprotective, and antidiabetic activities. Beyond its therapeutic relevance, *A. indica* has gained increasing attention in applied research for its roles in green nanotechnology, sustainable agriculture as a biopesticide, and environmental remediation via its bio-flocculant properties. The present review consolidates current evidence on the phytochemical composition and biological activities of *A. indica*, critically evaluates the mechanistic basis of its pharmacological actions, and identifies priority areas for future translational research and product standardisation.

**Citation:** Kapil, A., Anshul, A., Gupta, V., Barwal, S. K., & Singh, H. (2026). Phytochemical Diversity and Multispectral Pharmacological Activities of *Azadirachta indica* A.Juss. (Meliaceae): An Evidence-Based Review of Bioactive Constituents, Therapeutic Mechanisms, and Emerging Applications. *AgroEnvironmental Sustainability*, 4(2), 231–239. <https://doi.org/10.59983/s20260402013>

**Statement of Sustainability:** *Azadirachta indica* represents a sustainable botanical resource with significant potential to support environmental, agricultural, and public health goals. Its natural bioactive compounds offer eco-friendly alternatives to synthetic pesticides, antimicrobial agents, and chemical additives, thereby reducing ecological burdens associated with conventional products. Furthermore, neem-based applications in green nanotechnology, water treatment, and integrated pest management contribute to resource efficiency and environmental protection. Advancing evidence-based utilization, standardization, and conservation of *A. indica* can promote sustainable development while supporting biodiversity and community well-being.

## 1. Introduction

*Azadirachta indica* A.Juss., a member of the family Meliaceae, has occupied a central position in traditional medicinal systems of the Indian subcontinent for several millennia. Known colloquially as neem, the tree is referred to in Sanskrit literature as 'Sarva Roga Nivardini' (the curer of all ailments), a designation that reflects its extensive and varied ethnomedicinal applications in Ayurveda, Unani, Homeopathy, and Naturopathy (Reddy & Palagani, 2022). Contemporary pharmacological research has provided increasing experimental corroboration for many of these traditional uses, positioning neem among the most scientifically validated medicinal plants globally (Singh et al., 2019).

The renewed scientific interest in plant-derived bioactive compounds has been driven, in part, by escalating concerns regarding antimicrobial resistance and the adverse effects associated with synthetic pharmaceuticals. Phytotherapeutic agents generally offer



## ARTICLE HISTORY

Received: 21 January 2026

Revised: 19 February 2026

Accepted: 07 March 2026

Published: 15 June 2026

## KEYWORDS

antimicrobial activity  
antioxidant activity  
*Azadirachta indica*  
azadirachtin  
biopesticide  
neem  
phytochemistry

## COPYRIGHT

© 2026 Author(s)

e-ISSN: 2583-942X

Exclusive Rights: Society for  
AgroEnvironmental Sustainability,  
Dehradun, India

## LICENSE



This is an Open Access Article  
published under a CC BY 4.0  
International License

superior safety profiles and are frequently more accessible to populations in low- and middle-income countries, where approximately 80% of the population continues to rely on herbal medicine as the primary source of healthcare (Osman et al., 2025). Against this backdrop, *A. indica* represents a paradigmatic example of a medicinal plant whose phytochemical complexity has been translated into a diverse array of biological activities applicable across medicine, agriculture, and environmental science. Figure 1 shows multipurpose use of different plant parts such as leaves, bark, flowers, and seeds.




Figure 1. Multipurpose use of different plant parts such as leaves, bark, flowers, and seeds.

*A. indica* has been demonstrated to exert activity against both Gram-positive organisms, including *Staphylococcus aureus*, and Gram-negative pathogens such as *Salmonella dublin* and *Escherichia coli*, with inhibitory potency increasing proportionally with extract concentration (Osman et al., 2025). Additionally, the plant has been identified as a functional bio-flocculant in wastewater treatment owing to the positive surface charge of its extractable polymers, which facilitates the aggregation and removal of suspended pollutants (Zakaria et al., 2025). The purpose of the present review is to provide a comprehensive, evidence-based account of the phytochemical constituents and primary biological activities of *A. indica*, with particular attention to the mechanistic underpinnings of its pharmacological properties and its emerging translational applications.

## 2. Botanical Description

### 2.1. Taxonomic Classification

*A. indica* is classified within the order Sapindales, family Meliaceae, and is the sole species within the genus *Azadirachta* to achieve widespread botanical and pharmacological documentation (Table 1). The species was formally described by Antoine Laurent de Jussieu and its taxonomic status has remained largely stable in contemporary classification systems. From a cladistic perspective, the Meliaceae are distinguished by their production of limonoid terpenoids, of which azadirachtin in neem represents the most structurally complex and biologically significant example (Reddy & Palagani, 2022).



Table 1. Taxonomic classification of *A. indica*.

Taxonomic Rank	Classification
Kingdom	Plantae
Division	Magnoliophyta
Class	Magnoliopsida
Order	Sapindales
Family	Meliaceae
Genus	<i>Azadirachta</i> A.Juss.
Species	<i>Azadirachta indica</i> A.Juss.
Common Names	Neem (English); Nimtree; Margosa; Sarva Roga Nivarini (Sanskrit)

## 2.2. Morphological Characteristics

*A. indica* is a medium-to-large tropical evergreen tree attaining heights of 12–20 m under typical conditions and occasionally reaching 30 m in particularly favourable environments (Azad et al., 2023). Its broad, spreading crown and dense canopy of dark-green foliage render it highly valued as a shade tree throughout its natural and introduced range (Table 2). The leaves are pinnately compound, comprising 8–19 lanceolate leaflets with serrated margins, arranged alternately along the rachis (Bahri et al., 2025). The bark is characteristically thick, rough, and deeply fissured with a greyish-brown to reddish-brown colouration; it possesses a distinctly bitter taste attributable to its high concentration of triterpenoid constituents (Azis & Shofia, 2026).

The inflorescences are axillary panicles bearing small, white, bisexual flowers with a distinctive sweet fragrance, which appear predominantly during the spring season and are essential to the plant’s reproductive cycle (Nagendiran & Parimala, 2025). The fruit is a smooth, olive-like drupe measuring approximately 1–2 cm in length; it is green when immature and transitions to yellow upon ripening, each containing a single seed that serves as the primary commercial source of neem oil (Ajibade et al., 2026).

Table 2. Morphological characteristics and medicinal significance of plant parts of *A. indica*.

Plant Part	Morphological Description	Primary Medicinal Applications
Leaves	Pinnately compound; 8–19 lanceolate leaflets with serrated margins; dark green	Antimicrobial, antifungal, anti-inflammatory; treatment of skin diseases and febrile conditions
Bark	Thick, rough, fissured; greyish-brown to reddish-brown; intensely bitter taste	Antipyretic, antimalarial, anti-inflammatory; bark decoctions used in traditional fever management
Seeds	Enclosed in a single-seeded drupe; primary source of neem oil via cold-press extraction	Pesticidal, antifungal; source of azadirachtin and other limonoid insecticides
Flowers	Small, white, bisexual; arranged in axillary panicles; strongly fragrant	Traditional use in digestive disorders; anti-inflammatory activity reported
Roots	Lateral root system; bark similarly bitter to stem bark	Anthelmintic and antimicrobial applications in folk medicine
Oil (seed-derived)	Yellowish-brown fixed oil with characteristic odour; rich in fatty acids and limonoids	Pharmaceutical formulations, topical antimicrobials, cosmetics, agricultural pesticides

## 2.3. Geographical Distribution

*A. indica* is indigenous to the Indian subcontinent, where it grows naturally across diverse climatic zones ranging from semi-arid to sub-humid. It demonstrates exceptional adaptability, tolerating poor and degraded soils, extended drought, and high ambient temperatures, which has facilitated its naturalization in the broader tropical belt. Within South Asia, the species is widely distributed in Bangladesh, Nepal, Pakistan, and Sri Lanka (Reddy & Palagani, 2022). Beyond its native range, neem has been intentionally introduced into sub-Saharan Africa, Southeast Asia, Central America, and parts of the Caribbean and the Pacific, where it is cultivated for its shade, medicinal, and agricultural utility. The species’ drought resilience and ecological versatility make it one of the most cosmopolitan tropical medicinal trees.

## 3. Phytochemical Constituents

### 3.1. Classification of Bioactive Compounds

The phytochemical architecture of *A. indica* is exceptionally rich, encompassing several hundred identified compounds distributed across all anatomical parts of the plant (Chauhan et al., 2022). These constituents are conventionally categorised into two major chemical classes: isoprenoid (terpenoid) compounds and non-isoprenoid compounds. The former include a diverse array of limonoids most notably the tetranortriterpenoids as well as steroids and diterpenes (Table 3). The non-isoprenoid fraction comprises flavonoids, tannins, alkaloids, saponins, coumarins, phenolic acids, and their respective glycosides (Bahri et al., 2025; Ajibade et al., 2026; Azis & Shofia, 2026). The relative abundance and composition of these constituents vary considerably according to plant part, geographic origin, seasonality, and extraction methodology employed.



Among the major phytochemical classes: alkaloids confer antibacterial and antiprotozoal activities through mechanisms involving nucleic acid intercalation and membrane disruption; flavonoids function as potent radical scavengers and anti-inflammatory agents via inhibition of cyclooxygenase and lipoxygenase pathways; tannins exert astringent and antimicrobial effects through protein precipitation and membrane perturbation; saponins demonstrate immunomodulatory and surface-active properties; terpenoids the dominant class in neem contribute the principal insecticidal, antifeedant, and anti-inflammatory activities; and glycosides provide additional synergistic biological activity across multiple pharmacological targets (Bahri et al., 2025; Ajibade et al., 2026).

Table 3. Principal phytochemical constituents of *A. indica*: chemical class, distribution, and reported biological activity.

Compound / Class	Chemical Class	Primary Plant Part	Key Biological Activity	Selected Reference
Azadirachtin	Tetranortriterpenoid (Limonoid)	Seeds, leaves	Insecticidal, antifeedant, insect growth regulator; minimal toxicity to non-target organisms	Bahri et al. (2025); Nagendiran & Parimala (2025)
Nimbin	Triterpenoid (Limonoid)	Seeds, bark	Anti-inflammatory, antifungal, antipyretic	Bahri et al. (2025)
Nimbidin	Triterpenoid (Limonoid)	Seeds, bark	Antibacterial, anti-inflammatory, antiulcer	Bahri et al. (2025)
Salannin	Tetranortriterpenoid (Limonoid)	Seeds, leaves	Insect antifeedant; repellent activity against multiple pest species	Nagendiran & Parimala (2025)
Gedunin	Triterpenoid (Limonoid)	Seeds, bark	Antibacterial, antimalarial, antifungal	Nagendiran & Parimala (2025)
Quercetin	Flavonoid	Leaves, flowers	Antioxidant, anti-inflammatory, antiviral	Azis & Shofia (2026)
Rutin	Flavonoid glycoside	Leaves	Antioxidant, vascular protective, anti-inflammatory	Azis & Shofia (2026)
Catechins / Tannins	Polyphenol	Bark, leaves	Antimicrobial, astringent, antioxidant	Bahri et al. (2025)
Saponins	Triterpenoid glycoside	Leaves, seeds	Immunomodulatory, haemolytic, antimicrobial	Ajibade et al. (2026)
Alkaloids	Nitrogen-containing heterocycles	Leaves, bark	Antibacterial, antiprotozoal, analgesic	Bahri et al. (2025); Ajibade et al. (2026)
Terpenoids (general)	Isoprene-derived terpenes	All parts	Insecticidal, antimicrobial, anti-inflammatory	Ajibade et al. (2026)
Phenolic acids	Hydroxycinnamic / Hydroxybenzoic derivatives	Leaves, bark	Antioxidant, antimicrobial, anti-inflammatory	Azis & Shofia (2026)

### 3.2. Phytochemical Screening Studies

Systematic phytochemical screening of *A. indica* across multiple solvent systems and plant parts has consistently confirmed the presence of the compound classes enumerated above. Bahri et al. (2025) conducted methanol extraction of neem leaves and identified alkaloids, flavonoids, and tannins as major constituents via standard qualitative screening assays. Ajibade et al. (2026) characterised mixed distillates of neem alongside other medicinal plants, reporting the presence of terpenoids and glycosides as predominant classes. Azis and Shofia (2026) utilised ethanol-based extraction of neem leaf material and identified phenolics and flavonoids as the principal antioxidant-contributing fractions, corroborating findings reported by Nagendiran and Parimala (2025) using ethanolic flower extracts. These studies collectively affirm that solvent polarity substantially governs the extractable phytochemical profile, with polar solvents (methanol, ethanol, water) preferentially extracting flavonoids and phenolics, while less polar solvents enrich terpenoid and limonoid fractions.

## 4. Biological Activities

### 4.1. Antimicrobial Activity

The antimicrobial properties of *A. indica* extracts have been among the most thoroughly documented pharmacological attributes of the plant (Shah et al., 2021). Experimental investigations employing extracts from leaves, bark, and seeds prepared in various solvents have demonstrated inhibitory activity against a broad panel of bacterial and fungal pathogens (Table 4). Studies by Vadivel et al. (2025), Akabandi (2026), Tiranya and Munasinghe (2025), Azis and Shofia (2026), and Osman et al. (2025) collectively confirm significant antibacterial activity, particularly against *Staphylococcus aureus* and *Escherichia coli*, which serve as reference organisms for Gram-positive and Gram-negative antibacterial screening, respectively.

The mechanisms underlying the antimicrobial activity of neem extracts are multifactorial. Flavonoids and phenolic compounds are capable of forming complexes with bacterial cell wall proteins, thereby compromising structural integrity. Terpenoids, particularly limonoids such as nimbidin, have been shown to disrupt membrane permeability, leading to cellular lysis. Alkaloids may interca-



late with bacterial DNA and interfere with topoisomerase function. The broad-spectrum nature of neem's antibacterial action is therefore attributable to the synergistic action of multiple phytochemical classes acting on distinct molecular targets (Osman et al., 2025; Azis & Shofia, 2026).

Table 4. Summary of antimicrobial studies on *A. indica* extracts.

Author (Year)	Plant Part / Extract	Test Organism(s)	Principal Finding	Methodology
Vadivel et al. (2025)	Homeopathic dilutions of leaf	<i>S. aureus</i> , <i>E. coli</i>	Inhibitory activity demonstrated; correlated with LC-MS identified phenolic fractions	Disc diffusion, LC-MS analysis
Akabandi (2026)	Leaf extract (solvent unspecified)	Multiple bacterial strains	Antibacterial activity confirmed; inhibition zones recorded against reference strains	Disc diffusion / Agar well diffusion
Tiranya & Munasinghe (2025)	Whole plant extract	<i>E. coli</i> , <i>S. aureus</i>	Strong antibacterial activity; bioactive compound screening conducted concurrently	MIC determination, phytochemical screening
Azis & Shofia (2026)	Ethanol leaf extract	Skin-pathogenic bacterial and fungal strains	Broad-spectrum activity against dermatological pathogens	Disc diffusion, MIC/ MBC
Osman et al. (2025)	Aqueous bark extract	Multiple bacterial pathogens	Dose-dependent inhibition; activity increased proportionally with extract concentration	Agar well diffusion, spectrophotometric quantification

## 4.2. Antioxidant Activity

Oxidative stress arising from an imbalance between the generation of reactive oxygen species (ROS) and the capacity of endogenous antioxidant defence systems is implicated in the pathogenesis of numerous chronic diseases including cardiovascular disorders, diabetes mellitus, neurodegenerative conditions, and carcinogenesis. *A. indica* extracts have been shown to exhibit potent free-radical scavenging activity in multiple in vitro assay systems, including DPPH (2,2-diphenyl-1-picrylhydrazyl) radical scavenging and ferric-reducing antioxidant power (FRAP) assays (Suthar et al., 2025; Nagendiran & Parimala, 2025; Yahaya et al., 2025). The principal antioxidant-active constituents are the leaf phenolics particularly flavonoids such as quercetin and rutin which function as hydrogen donors capable of neutralising ROS through electron transfer and hydrogen atom transfer mechanisms.

Yahaya et al. (2025) reported comparative in vitro antioxidant activity of n-hexane and ethyl acetate fractions of *A. indica*, demonstrating fraction-dependent variation in radical scavenging capacity that reflects the differential distribution of active phenolic constituents. Nagendiran and Parimala (2025) similarly documented antioxidant activity in ethanolic flower extracts, expanding the range of plant parts characterised for antioxidant function. These findings collectively support the utility of neem-derived phytochemicals in the development of natural antioxidant formulations applicable in both nutraceutical and pharmaceutical contexts.

## 4.3. Immunomodulatory Activity

The immunomodulatory properties of *A. indica* have been evaluated in both monotherapy and combination therapy experimental models. Suthar et al. (2025) demonstrated that the administration of *A. indica* extract, both alone and in combination with *Tinospora cordifolia*, produced significant enhancement of antioxidant and immunological parameters in broiler chickens, indicating synergistic augmentation of host immune responses when neem is used in conjunction with other immunomodulatory botanicals. The immunostimulatory effects of neem are partly attributed to its saponin constituents, which have been demonstrated to stimulate lymphocyte proliferation and macrophage activation, as well as to its polysaccharide fractions that may function as adjuvant-like immunopotentiators.

## 4.4. Neuroprotective Activity

Emerging evidence from preclinical investigations points to a neuroprotective role for *A. indica* phytochemicals in models of neurological dysfunction. Rao et al. (2025) demonstrated that neem extract administration ameliorated memory deficits and reduced anxiety-like behaviours in an animal model of depression, with the observed effects attributed to modulation of cholinergic neurotransmission. The cholinergic hypothesis of cognitive decline proposes that impairment of acetylcholinergic signalling in the hippocampus and cortex underlies deficits in learning and memory; neem extracts may therefore exert their cognitive protective effects through inhibition of acetylcholinesterase or upstream neuroprotective mechanisms involving anti-inflammatory and antioxidant pathways. These findings justify further mechanistic investigation and eventual clinical evaluation of neem-derived compounds for the management of neurodegenerative disorders.



## 4.5. Antidiabetic Activity

Multiple bioactive compounds present in *A. indica* have demonstrated potential in modulating glucose homeostasis and metabolic parameters relevant to diabetes mellitus (Table 5). Nagendiran and Parimala (2025) conducted comparative evaluation of antidiabetic activity in ethanolic extracts of *A. indica* and *Melia dubia* flowers, reporting significant alpha-amylase and alpha-glucosidase inhibitory activity that suggests interference with post-prandial glucose absorption at the intestinal level. The antidiabetic mechanisms proposed for neem phytochemicals include: inhibition of carbohydrate-hydrolysing enzymes (alpha-amylase, alpha-glucosidase); enhancement of peripheral glucose uptake via insulin-sensitising effects; reduction of oxidative stress in pancreatic beta cells; and modulation of lipid metabolism, which is often dysregulated in type 2 diabetes. The flavonoid quercetin, present in significant concentrations in neem leaf, has been independently documented as an inhibitor of protein tyrosine phosphatase 1B, a validated antidiabetic drug target (Nagendiran & Parimala, 2025).

Table 5. Consolidated summary of pharmacological activities of *A. indica*.

Pharmacological Activity	Active Constituent(s) Implicated	Proposed Mechanism(s)	Key Reference(s)
Antimicrobial	Nimbidin, alkaloids, tannins, phenolics	Cell wall disruption, membrane permeabilisation, enzyme inhibition, DNA intercalation	Vadivel et al. (2025); Osman et al. (2025); Azis & Shofia (2026)
Antioxidant	Quercetin, rutin, flavonoids, phenolic acids	Free-radical scavenging via hydrogen atom transfer and electron transfer mechanisms; ROS neutralisation	Suthar et al. (2025); Nagendiran & Parimala, (2025); Yahaya et al. (2025)
Immunomodulatory	Saponins, polysaccharides, polyphenols	Lymphocyte proliferation stimulation; macrophage activation; cytokine modulation	Suthar et al. (2025)
Neuroprotective	Terpenoids, flavonoids, phenolics	Acetylcholinesterase inhibition; anti-inflammatory neuroprotection; antioxidant-mediated neuronal protection	Rao et al. (2025)
Antidiabetic	Quercetin, flavonoids, terpenoids	Alpha-amylase and alpha-glucosidase inhibition; insulin sensitisation; pancreatic beta-cell protection	Nagendiran & Parimala (2025)
Insecticidal / Biopesticidal	Azadirachtin, salannin, nimbin	Ecdysteroid antagonism; antifeedant activity; hormonal disruption of insect moulting cycle	Cece et al. (2026); Bahri et al. (2025)
Bio-flocculant / Water Treatment	Cationic polymers, phenolics	Charge neutralisation of suspended particulates; floc formation and sedimentation enhancement	Zakaria et al. (2025)

## 5. Nanotechnology Applications

### 5.1 Green Biosynthesis of Nanoparticles

The application of plant extracts in the environmentally sustainable synthesis of metal nanoparticles represents an active frontier of nanoscience research. *A. indica* extracts have been extensively investigated for their capacity to function as reducing and stabilising agents in the synthesis of silver and iron nanoparticles, offering a non-toxic, energy-efficient alternative to conventional chemical reduction methods (Siddiqui et al., 2026). The flavonoids, terpenoids, and phenolic compounds present in neem extracts are principally responsible for the reduction of ionic silver ( $Ag^+$ ) and ferric iron ( $Fe^{3+}$ ) to their respective zero-valent metallic nanoparticle forms, while simultaneously capping the nascent nanoparticles to prevent aggregation and confer colloidal stability.

### 5.2. Biomedical Applications of Neem-Mediated Nanoparticles

Silver nanoparticles synthesised via neem-mediated green routes have demonstrated superior antimicrobial activity against a broad spectrum of pathogenic microorganisms in comparison with unmodified plant extracts alone. The enhanced activity is attributable to the high surface area-to-volume ratio of nanoparticles, which facilitates intimate contact with microbial cell walls and membranes, promoting oxidative damage and cellular dysfunction (Siddiqui et al., 2026). Furthermore, neem-mediated nanoparticles have been investigated for catalytic applications in environmental remediation, capitalising on their high surface reactivity to accelerate the degradation of chemical pollutants. These attributes position *A. indica*-mediated nanoparticles as multifunctional agents at the interface of green chemistry, nanomedicine, and environmental biotechnology.

## 6. Agricultural Applications



## 6.1. Biopesticidal Properties

The biopesticidal utility of *A. indica* is well established and represents one of its most commercially developed applications. Azadirachtin, the principal active compound, exerts its insecticidal effects primarily through antagonism of ecdysone, the principal insect moulting hormone, thereby disrupting larval development, ecdysis, and reproductive function. This mechanism of action, which is highly specific to arthropods and does not affect vertebrate hormonal systems, confers upon azadirachtin an excellent selectivity profile relative to conventional broad-spectrum synthetic insecticides (Cece et al., 2026). Additional limonoids salannin, nimbin, and nimbidin contribute antifeedant and repellent activities, further reducing pest fitness and population pressure.

## 6.2. Efficacy Against Economically Important Pests

Cece et al. (2026) evaluated the biopesticidal activity of *A. indica* seed and leaf extracts against the fall armyworm (*Spodoptera frugiperda*), one of the most economically damaging polyphagous pests affecting maize production globally. The investigators documented significant inhibition of larval feeding behaviour and reductions in survival rates, supporting the utility of neem-based formulations as integrated pest management tools in sub-Saharan African agricultural systems. The ecological safety profile of neem-based pesticides, arising from their rapid photodegradation and low persistence in soil, reduces the risk of non-target toxicity and environmental bioaccumulation compared with synthetic organophosphate and pyrethroid alternatives.

## 7. Environmental Applications

### 7.1. Wastewater Treatment and Bio-flocculant Properties

The application of *A. indica*-derived materials in water treatment is an emerging area of environmental biotechnology. The presence of cationic polymers and polyphenolic compounds in neem extracts enables these materials to function as natural coagulants and bio-flocculants, aggregating suspended particulate matter through charge neutralisation and bridging mechanisms. Zakaria et al. (2025) demonstrated the efficacy of *A. indica* as a bio-flocculant in reducing turbidity and suspended solids in landfill leachate, a particularly recalcitrant wastewater stream characterised by high pollutant load and variable composition. The performance of neem-based bio-flocculants was attributed to the positive surface charge of extractable components, which effectively neutralises the negative charge of colloidal particles in suspension, inducing floc formation and subsequent sedimentation.

The use of plant-derived coagulants in water treatment offers several sustainability advantages over inorganic chemical coagulants such as aluminium sulphate: they are biodegradable, produce lower sludge volumes, do not introduce toxic metal residues into treated water, and can be sourced from locally available plant biomass in low-income settings (Zakaria et al., 2025). These characteristics position *A. indica* as a candidate for incorporation into sustainable, decentralised water treatment systems, particularly in tropical regions where the plant grows abundantly.

## 8. Future Research Perspectives

Despite the considerable body of evidence accumulated to date, several critical knowledge gaps remain that must be addressed to realise the full translational potential of *A. indica*. At the phytochemical level, systematic isolation, structural elucidation, and quantitative profiling of minor constituent classes particularly the non-limonoid alkaloids and phenylpropanoids are needed across a broader range of geographic accessions and seasonal time points. Advances in high-resolution mass spectrometry and nuclear magnetic resonance-based metabolomics offer powerful tools for comprehensive phytochemical characterisation that should be applied systematically to this species.

In the pharmacological domain, the majority of available evidence derives from *in vitro* assay systems and limited animal model studies; rigorous randomised controlled clinical trials evaluating the efficacy and safety of standardised neem formulations in human subjects are conspicuously absent for most claimed indications. Mechanistic studies employing molecular docking, target identification, and pharmacokinetic profiling are required to validate proposed mechanisms of action and to identify the most promising lead compounds for drug development pipelines (Suthar et al., 2025; Rao et al., 2025).

In the domain of agricultural applications, field-scale evaluation of neem-based biopesticide formulations under diverse agroclimatic conditions, and their integration into evidence-based integrated pest management protocols, represents a priority research area (Cece et al., 2026). Standardisation of neem-based products, encompassing phytochemical authentication, quality control, and stability testing, is prerequisite for regulatory approval and commercialisation of herbal formulations in both domestic and international markets. The application of nanotechnology to enhance the bioavailability and targeted delivery of neem-derived phytochemicals likewise represents a productive intersection of multiple research frontiers (Siddiqui et al., 2026).

## 9. Conclusion

*A. indica* is a botanically and pharmacologically exceptional species whose extensive phytochemical diversity underpins a correspondingly broad spectrum of experimentally validated biological activities. The present review has synthesised evidence demonstrating that neem extracts, derived from multiple anatomical parts of the plant, exhibit antimicrobial, antioxidant, immunomodulatory, neuroprotective, and antidiabetic activities attributable to a complex and synergistic ensemble of limonoid terpenoids,



flavonoids, tannins, alkaloids, saponins, and phenolic acids. Beyond therapeutic applications, *A. indica* has demonstrated substantial utility in green nanotechnology, as a source of sustainable biopesticides in integrated crop protection, and as a bio-flocculant in environmentally responsible water treatment processes. The convergence of traditional ethnomedicinal knowledge with contemporary experimental investigation positions *A. indica* as a priority candidate for translational pharmacognosy. Future research should prioritise: rigorous clinical evaluation of standardised neem formulations; advanced mechanistic studies elucidating the molecular basis of pharmacological activity; comprehensive phytochemical profiling using modern analytical platforms; and systematic quality standardisation of neem-derived products for regulatory acceptance. Interdisciplinary collaboration spanning phytochemistry, pharmacology, toxicology, agronomy, and environmental science will be essential to translate the accumulated basic science evidence into tangible therapeutic, agricultural, and environmental innovations.

### Author Contributions

Archit Kapil: Conceptualization; Data curation; Investigation; Methodology; Resources; Software; Validation; Visualization; Writing – original draft; Writing – review and editing; Anshul: Data curation; Methodology; Software; Visualization; Writing – original draft; Vansh Gupta: Data curation; Methodology; Visualization; Writing – original draft; Sandeep Kumar Barwal: Supervision; Writing – original draft; Writing – review and editing; Harsh Singh: Writing – original draft; Writing – review and editing.

### Funding Information

This research received no external funding.

### Acknowledgements

The authors acknowledge the School of Botany, Maa Shakumbhari University, Saharanpur, for institutional support throughout the preparation of this manuscript.

### Declarations

**Conflicts of interest:** The authors report no conflict of interest in relation to the publication of this manuscript.

**Institutional/Ethics approval:** As this study did not involve human or animal subjects, ethical approval was not required.

**Data availability:** The data supporting the conclusions of this study can be obtained from the corresponding author upon reasonable request.

**Supplementary material:** No supplementary material is associated with this manuscript.

### References

- Ajibade, G., Abdulazeez, R., Abdullahi, I., Achi, M., & Nuhu, A. (2026). Preliminary phytochemical screening and characterization of *Artemisia annua*, *Azadirachta indica*, *Nauclea latifolia* and *Vernonia ambigua* distillates. *Kasu Journal of Chemical Science*, 1(2), 1–11.
- Akabandi, J. (2026). *Antibacterial activity of leaf extract of Azadirachta indica against selected bacteria* [Master's thesis, C.K. Tedam University of Technology and Applied Sciences]. <https://doi.org/10.13140/RG.2.2.28325.36322>
- Azad, M. A. K., Barwal, S. K., & Moni, A. (2023). Exploring the impact of integrated breeding strategies in enhancing yield, nutritional quality, and stress tolerance in alfalfa. *Plant Trends*, 1(1), 1–15.
- Azis, F., & Shofia, V. (2026). Phytochemical evaluation and antimicrobial activity of neem leaf (*Azadirachta indica* A. Juss.) extract against skin pathogens. *Medical Sains: Jurnal Ilmiah Kefarmasian*, 11, 1–6. <https://doi.org/10.37874/ms.v11i1.1803>
- Bahri, S., Hijriani, B., & Widullutfi, W. (2025). Phytochemical screening of methanol extract of neem (*Azadirachta indica*) leaves. *Jurnal Biologi Tropis*, 25, 6115–6121. <https://doi.org/10.29303/jbt.v25i4.10998>
- Cece, M., Olayemi, I., Salaudeen, M., Ukubuiwe, A., & Salihu, I. (2026). Biopesticidal activity of seed and leaf extracts of *Azadirachta indica* and *Jatropha curcas* against fall armyworm (*Spodoptera frugiperda*) in maize. *Journal of Entomology and Agronomy Studies*, 10, 1–11. <https://doi.org/10.70382/bjeas.v10i1.023>
- Chauhan, C., Goutam, C., Kumar, S., Singh, I., & Yerramilli, V. (2022). GC-MS analysis and phytochemical screening of chloroform extract of *Amaranthus viridis*. *The Journal of the Indian Botanical Society*, 102(01), 53–60.
- Nagendiran, S., & Parimala. (2025). Comparative study on phytochemical screening, antioxidant, antimicrobial and antidiabetic activity in the ethanolic extract of *Azadirachta indica* and *Melia dubia* flowers. *International Journal of Zoological Investigations*, 11, 1017–1024. <https://doi.org/10.33745/ijzi.2025.v11i02.100>
- Osman, W., Gibreel, H., Rabia, R., Salih, M., Mirghani, M., Ashour, A., & Waggiallah, H. (2025). Antimicrobial effects of aqueous bark extracts of *Khaya senegalensis*, *Azadirachta indica* and *Eucalyptus camaldulensis* against bacterial pathogens. *Salud, Ciencia y Tecnología*, 6, 2683. <https://doi.org/10.56294/saludcyt20262683>
- Rao, B., Raza, S., & Tiwari, P. (2025). *Azadirachta indica* ameliorates memory deficits and reduces anxiety-like behavior by modulating cholinergic neurotransmission in an animal model of depression. *Traditional and Integrative Medicine*, 10(3), 19758. <https://doi.org/10.18502/tim.v10i3.19758>
- Reddy, I. V. S., & Palagani, N. (2022). Neem (*Azadirachta indica*): A review on medicinal Kalpavriksha. *International Journal of Economic Plants*, 9, 59–63. <https://doi.org/10.23910/2/2021.0437d>
- Shah, S. H., Kumar, S., & Yerramilli, V. (2021). An insight on the ascorbate peroxidase and glutathione reductase activities in plants under salinity stress: Mini review. *Research Journal of Agricultural Sciences*, 12, 1558–1563.



- Siddiqui, A., Fatima, N., Yadav, A., Sinha, A., Tripathi, V., Singh, G., Kumar, A., & Bhadauria, V. (2026). Eco-friendly biosynthesis of recyclable silver nanocatalyst and iron nanocatalyst using *Azadirachta indica*. *YMER Digital*, 24.
- Singh, T. B., Kumar, S., Singh, D. K., Khirwar, S. S., & Agnihotri, R. K. (2019). Toxicity of Cr and Pb during vegetative growth of *Sesamum indicum* L. *The Journal of Indian Botanical Society*, 98(3–4), 219–226.
- Suthar, P., Sadariya, K., Vaghela, S., & Bhavsar, S. (2025). Evaluation of immunomodulatory and antioxidant activities of *Azadirachta indica* and *Tinospora cordifolia* alone and in combination in broilers. *Journal of Phytopharmacology*, 14, 393–404. <https://doi.org/10.31254/phyto.2025.14514>
- Tiranya, A., & Munasinghe, M. (2025). Screening of bioactive compounds and antibacterial efficacy of *Azadirachta indica* extract against *Escherichia coli* and *Staphylococcus aureus*. *Conference Proceeding: Medicinal Plants, December 18–20, 2025, Sri Lanka*.
- Vadivel, K., Singh, S., Paul, S., Shaw, N., & Paul, S. (2025). Phytochemical analysis, LC-MS analysis and antimicrobial activity of homeopathic dilutions of *Azadirachta indica* against *Staphylococcus aureus* and *Escherichia coli*. *Journal of Homeopathy*, 5(2), 1–7.
- Yahaya, A., Musa Bola, B., Yusuf, S., & Yunusa, I. (2025). Comparative in vitro antioxidant and trypanocidal activities of n-hexane and ethyl acetate extracts of *Azadirachta indica* on *Trypanosoma brucei*. *Conference Proceeding: Sivas Cumhuriyet University: 4th International Congress of Veterinary Biochemistry and Clinical Biochemistry*.
- Zakaria, S. N. F., Zakaria, F., Selvam, M., Peter, E., Aziz, H., Mohamad, M., & Bolong, N. (2025). Performance of *Azadirachta indica* as bio-flocculant in reducing turbidity concentration from landfill leachate. *International Journal of Integrated Engineering*, 17, 194–201. <https://doi.org/10.30880/iji.e.2025.17.03.021>

**Publisher's note/Disclaimer:** Regarding jurisdictional assertions in published maps and institutional affiliations, We maintains its neutral position. All publications' statements, opinions, and information are the sole responsibility of their respective author(s) and contributor(s), not We or the editor(s). We and/or the editor(s) expressly disclaim liability for any harm to persons or property caused by the use of any ideas, methodologies, or products described in the content.